

The **fluTAS report** is a fortnightly influenza season update produced by the Communicable Disease Prevention Unit to inform healthcare organisations and the public about the current level of influenza disease activity in Tasmania. Multiple data sources are used to obtain some measure of influenza disease activity in the community.


Summary

There has been an increase in Tasmanian influenza notifications during the last fortnight. Syndromic surveillance systems are reporting increasing levels of influenza like illness (ILI) above the baseline. Respiratory pathogen testing suggests that influenza and other non-influenza viruses are contributing to ILI in Tasmania.

Influenza notifications

Tasmanian laboratories are required to notify evidence of influenza infection in specimens collected from patients. These specimens are usually nose or throat swabs but some may be blood tests.

Influenza notifications 2012 - 2013 (at 1 September 2013)



There has been an increase in the number of influenza notifications, with 42 cases received during the last fortnight ending 1 September. These notifications account for 45% of the 93 influenza notifications now received during 2013.

The majority of notifications received have been in residents of the South region of Tasmania. Differences in notifications for the year to date may not reflect the different level of influenza between regions. Apart from differences in regional populations, this is also affected by how many people seek medical care and get tested for influenza.


Location of influenza notifications 2013 (at 1 September 2013)

	SOUTH	NORTH	NORTH WEST	TOTAL
Notifications	70	15	8	93

Laboratory testing

Influenza types

During 2013 up to 1 September, 78 Influenza A notifications and 15 Influenza B notifications were received. The majority of typed Influenza A cases have been Influenza A H1N1 (2009 strain).



Influenza type in confirmed cases 2013 (Notifications Received at 1 September 2013)

	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	TOTAL
Influenza A	2	4	2	3	2	3	12	50					78
Influenza B	1	0	0	2	0	0	4	8					15


A wide range of infectious germs (mostly viruses) commonly cause winter coughs, colds and influenza-like illnesses. Some people with these symptoms will visit their doctor. The decision whether to test someone for influenza rests with their treating doctor, and depends on their symptoms. The best test for influenza is a PCR test, which detects influenza virus RNA in a nose or throat swab. The number of these tests being performed in public and private Tasmanian laboratories is a useful indicator of the level of respiratory illness in the community.

The total number of weekly influenza PCR tests performed in Tasmania has remained at an increased level over recent weeks. In the last fortnight 16% of PCR tests performed were positive for influenza.

Respiratory pathogen types

The Royal Hobart Hospital (RHH) performs additional PCR tests on nose and throat swabs to detect other non-influenza respiratory pathogens which cause illness. These specimens have been mostly collected from state-wide Emergency Departments or hospitalised patients. The level of PCR testing by the RHH Laboratory remained steady during the previous fortnight, as did the number of positive results reported. The proportion of tests detecting Influenza A increased. While Rhinovirus was most frequently detected, the proportion of tests detecting Respiratory Syncytial Virus (RSV) decreased.

RHH Respiratory Pathogen Testing 2013 (1-Jan to 1-Sep)



Influenza-like illnesses (Syndromic Surveillance)


Influenza-like illness (ILI) is much more common than microbiologically confirmed influenza. For much of the year, common colds and other respiratory illnesses make up most of the ILI occurring in the community. However, during the annual influenza season, the proportion of the population experiencing symptoms of ILI who have influenza usually increases. It is therefore useful to monitor the proportion of people reporting ILI, regardless of the cause.

FluTracking

FluTracking is a weekly online survey that asks participants to report whether they have had fever and cough in the preceding week. It is a joint initiative of Newcastle University, Hunter New England Population Health and the Hunter Medical Research Institute. For more information, go to www.flutracking.net.

An increasing proportion of Tasmanian FluTracking participants reported ILI symptoms during the most recent fortnight ending 1 September. Unvaccinated participants reported a higher level of ILI than vaccinated participants.

FluTracking - Tasmanian participants 2012 - 2013 (at 1-Sep)



General practice surveillance

ASPREN is a network of registered sentinel GPs throughout the state who report fortnightly on the number and proportion of presentations of patients with fever, cough and fatigue. ASPREN is a joint initiative of the Royal Australian College of General Practitioners and University of Adelaide: www.dmac.adelaide.edu.au/aspren.

Tasmanian data from participating General Practices up to the fortnight ending 25 August 2013 indicated a fluctuating level of influenza-like illness presentations recently but at levels lower than reported at this time during 2012.

Other measures of influenza activity

FluCAN

The Influenza Complications Alert Network (FluCAN) reports on influenza related hospitalisations and complications in sentinel hospitals in each state including Tasmania. Over the fortnight ending 2 September 2013 the number of people with influenza admitted to participating sentinel hospitals continued to increase.

Interstate activity

The Australian Influenza Report is compiled from a number of data sources, including laboratory-confirmed notifications to NNDSS, sentinel influenza-like illness reporting from general practitioners and emergency departments, workplace absenteeism, and laboratory testing. The current national report is available at <http://www.health.gov.au/internet/main/publishing.nsf/content/cda-surveil-ozflu-flucurr.htm>.

There has been a steady increase in influenza activity in most Australian States and Territories. Nationally the majority of recent notifications have been due to Influenza A. At the State and Territory level there has been variation as to whether Influenza A H1N1, Influenza A H3N2 or Influenza B is the major strain circulating in that jurisdiction.

Annual Influenza Vaccine

The contents of the annual influenza vaccine are reviewed late each year, aiming to have vaccines produced for the following year that provide protection from the strains of influenza that are likely to be common during winter. The recommended formulation of the 2013 vaccine includes two significant changes from the vaccines used in 2011 and 2012. These are described at <http://www.tga.gov.au/about/committees-aivc.htm>.

Annual vaccination is recommended and is free* for Tasmanians at risk of severe influenza, including:

- anyone aged 65 and over
- Indigenous people who are aged 15 years or over
- pregnant women
- any person six months of age and over with a chronic condition predisposing to severe influenza illness that requires regular medical follow-up or hospitalisation such as: cardiac disease, respiratory disease including severe asthmatics, kidney disease, diabetes, impaired immunity, neuromuscular disease.

* The cost of the vaccine is covered for these groups; there may be a consultation fee for the medical provider to administer the vaccine.

Help prevent the spread of flu and other respiratory viruses

Flu and other respiratory viruses are easily passed from person to person through the air when infected people cough or sneeze without covering their mouths or noses. As well as getting the seasonal flu vaccination, there are some basic things you can do to help prevent the spread of flu and other respiratory viruses.



Wash your hands regularly, especially before touching your face, food or utensils.



If you do get sick, please don't share it – cover your mouth when you cough or sneeze with the inside of your elbow. If you cover a cough or sneeze with your hands, remember to wash your hands afterwards.



Stay away from work or school if you are ill.



The **fluTAS Report** is a fortnightly flu season update produced by the Public and Environmental Health Service to inform healthcare organisations and the public about the current level of flu activity in Tasmania.

Alongside routine surveillance of diseases in Tasmania, the report combines multiple data sources to obtain a measure of flu activity in the community, which can be used by our health system to prepare and respond.

To provide feedback on the fluTAS Report, email the [Communicable Disease Prevention Unit](#) or call the Public Health Hotline on 1800 671 738.